

ANNEX A

Percentage changes from BTO and RSPB surveys, for the 8 species showing large declines (over 25% declines) according to both surveys, (plus hawfinch and redstart, which show less large declines across both sets of data, but where RSPB coverage was more effective); and for the 11 species showing large increases (over 25% increases) according to both surveys.

Species	BTO survey sites (% change)	RSPB survey sites (% change)
Declining species		
Lesser redpoll	-88.9	-58.7
Willow tit	-77.5	-72.5
Willow warbler	-74.2	-68.8
Spotted flycatcher	-70.4	-36.3
Tree pipit	-69.7	-85.4
Wood warbler	-64.0	-55.0
Lesser spotted woodpecker	-43.6	-58.9
Garden warbler	-25.6	-39.4
<i>Hawfinch*</i>	-17.4	-73.5
<i>Redstart**</i>	7.7	-54.4
Increasing species		
Chiffchaff	+154.8	+190.7
Goldcrest	+138.3	+87.5
Green woodpecker	+80.7	+269.3
Coal tit	+74.0	+48.7
Great spotted woodpecker	+69.8	+123.1
Robin	+63.5	+71.3
Blackcap	+57.2	+79.8
Wren	+56.5	+91.0
Treecreeper	+51.5	+95.1
Great tit	+51.2	+31.8
Blue tit	+30.8	+32.5

*BTO sample of hawfinch was very small.

**RSPB surveys covered more of this species core range.

Other species with more moderate results:

Blackbird	+15.8%	+64.3%
Bullfinch	+10.7%	-1.9%
Chaffinch	+25.9%	-5.5%
Dunnock	-5.8%	+13.0%
Jackdaw	+6.9%	-19%
Jay	-19.9%	-26.8%
Long tailed tit	+130.8%	+12.7%
Marsh tit	-27.0%	+ 26.5%
Mistle Thrush	-12.5%	+8.8%
Nuthatch	+78.7%	+9.8%
Pied flycatcher	-24.6%	-20.1%
Siskin	+107.6%	+28.4%
Songthrush	+15.7%	+52.2%
Woodpigeon	N/a	+85.5%