

ForestGALES

A PC-based wind risk model
for British Forests

User's Guide

Version 2.1 2006

Barry Gardiner, Juan Suárez,
Alexis Achim, Sophie Hale
and Bruce Nicoll

© Crown Copyright 2006

ISBN: 0 85538 632 0

Produced in the United Kingdom
FCSW001/FG(KMA)/AST&AM-200/DECO6

Licence Agreement

This licence agreement applies to ForestGALES version 2.1

You should carefully read the following terms and conditions before using this software. Your use of this software indicates your acceptance of this licence agreement and warranty.

You shall not alter, modify or reverse engineer the original software or give anyone else permission to do so. Unregistered use of ForestGALES after the beta-testing evaluation period will be in violation of UK and international copyright laws.

Disclaimer of Warranty

THIS SOFTWARE AND THE ACCOMPANYING FILES ARE DISTRIBUTED 'AS IS' AND WITHOUT WARRANTIES AS TO PERFORMANCE OR MERCHANTABILITY OR ANY OTHER WARRANTIES WHETHER EXPRESSED OR IMPLIED. In particular, there is no warranty for the predictions made by the model as they are regarded as indicative and not prescriptive.

NO RESPONSIBILITY FOR LOSS OCCASIONED TO ANY PERSON OR ORGANISATION ACTING, OR REFRAINING FROM ACTION, AS A RESULT OF ANY MATERIAL IN THIS PRODUCT CAN BE ACCEPTED BY THE FORESTRY COMMISSION.

ForestGALES is a PC-based program that has been designed to work under all Microsoft Windows Operating Systems. Because of the various hardware environments into which ForestGALES may be put, NO WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE IS OFFERED.

The installation disks have been thoroughly tested for computer viruses with the latest virus checking technologies available at the Forestry Commission. However, the user must assume the entire risk of using the ForestGALES program and therefore nobody apart from the user will be responsible for the loss of critical data or any other damage to the systems as a result of virus attack or a malfunction of the program.

Trademarks

All terms mentioned in this book that are known to be trademarks or service marks have been appropriately indicated. The Forestry Commission cannot attest to the accuracy of this information. Use of a term in this book should not be regarded as affecting the validity of any trademark or service mark. Readers should contact the appropriate companies for more complete information regarding trademarks and registration.

	Preface	v
1.	Introduction	1.1
	Why use ForestGALES?	1.1
	The degree of constraint posed by strong winds in Britain	1.1
	Historical context – Previous predictive windthrow model	1.1
	What does ForestGALES calculate?	1.2
	How does ForestGALES compare to the WHC?	1.2
	What is new in ForestGALES 2	1.3
	Wind climate	1.3
	Appearance	1.3
	Defaults	1.3
	Multiple stands (batch) mode	1.3
	Export	1.4
2.	Installing and uninstalling	2.1
	System requirements	2.1
	Installing ForestGALES	2.1
	Uninstalling ForestGALES	2.2
	Running the model for the first time	2.3
3.	How the model works	3.1
	How does the model work?	3.1
	What force would be needed to uproot or break the tree?	3.1
	What wind speed would create the force required to damage the tree?	3.1
	What is the probability of a damaging wind speed occurring?	3.2
4.	Using ForestGALES	4.1
	Running ForestGALES	4.1
	Operating modes	4.1
	Selecting a mode and opening a query form	4.1
	ForestGALES query forms	4.1
	Navigating ForestGALES	4.3
	Using the mouse to make selections	4.3
	Using the Tab key	4.3
	Using shortcuts	4.4
	Getting help	4.4
5.	Predictions using field measurements	5.1
	Introduction	5.1
	The stand characteristics box	5.1
	Options – Stand ID	5.1
	Options – Cultivation	5.1
	Options – Drainage	5.2
	Options – Soil type	5.2
	Options – Current spacing	5.2
	Options – Current stocking	5.2
	The tree characteristics box	5.3
	Options – Species	5.3
	Options – Top height	5.3
	Options – DBH	5.4
	The DAMS box	5.4
	Grid references method	5.4
	Exact calculation method	5.4
	Rough guess method	5.5
	The upwind edge effect box	5.5

Contents

	The controls box	5.6
	The wind damage risk box	5.6
	Critical wind speed	5.6
	Return period	5.6
	Wind damage risk status	5.7
	Windthrow Hazard Classification (WHC)	5.7
6.	Predictions using yield models	6.1
	Introduction	6.1
	Stand characteristics box	6.1
	Options – Planting year	6.1
	Tree characteristics box	6.2
	Options – Yield class	6.2
	Options – Thinning regime	6.2
	Options – Initial spacing	6.2
	Options – Age	6.2
	The 'Tree Details' button	6.2
	Other boxes	6.2
	User defined yield tables	6.3
	Overview	6.3
	Naming user defined yield models	6.3
	Using Excel® to create user defined yield models	6.4
7.	Predictions through time	7.1
	Introduction	7.1
	Stand characteristics box	7.1
	Tree characteristics box	7.1
	The DAMS box	7.1
	The controls box	7.1
	User-defined yield tables	7.1
	Viewing the results	7.2
	The graphics window	7.2
	The tabular data windows	7.3
	The results form control box	7.3
8.	Multiple stands	8.1
	Introduction	8.1
	The multiple stands helper	8.2
	The controls box	8.3
	The multiple stands query form	8.3
	The controls box	8.4
	Year of calculation	8.4
9.	Examples	9.1
	Single stand predictions using field measurements – getting started	9.1
	Single stand predictions using field measurements – the effect of a new edge	9.1
	Single stand predictions using yield models	9.2
	DAMS example	9.3
	DAMS – rough guess example	9.4
	DAMS – grid reference example	9.4
	DAMS – calculate DAMS example	9.5
10.	Glossary	10.1
11.	References & further reading	11.1

Welcome to ForestGALES

Acknowledgements

The initial development of ForestGALES was funded by the Forestry Commission, and in part by the European Union as part of the STORMS project. Continued improvement has been supported by the Forestry Commission (Corporate and Forestry Support). We are grateful to the many users of ForestGALES who have provided valuable feedback and comments on the previous version. We would also like to thank all those who tested the Beta version of this software. Colleagues within the Forestry Commission who have contributed to the development include Bill Rayner, Bill Mason, Colin Edwards, Elaine Dick, Kirstie Adamson, Duncan Ray, Shaun Mochan and Chris Quine. We would like to thank all those who have helped make this project a success.

Credits

Stability project team:

Barry Gardiner, Bruce Nicoll, Juan Suárez, Alexis Achim, Sophie Hale, Elspeth Macdonald and Bill Rayner

Software development:

Juan Suárez, Barry Gardiner, Roger Dunham, Juergen Bohl and Mark Brady

Testing and validation:

Juan Suárez, Julie Barrette and Sophie Hale

Manual:

Barry Gardiner, Juan Suárez, Alexis Achim, Sophie Hale and Bruce Nicoll.

For details of latest updates and on-line version of this manual access

www.forestresearch.gov.uk/forestgales

For help and assistance contact:

Bruce Nicoll
Forest Research
Northern Research Station
Roslin
Midlothian
EH25 9SY
United Kingdom

Phone: + 44 131 445 2176

Fax: + 44 131 445 5124

E-mail: forestgales.support@forestry.gsi.gov.uk



Why use ForestGALES?

Wind damage to forest stands results in both direct costs (additional cost of harvesting) and indirect costs (loss of amenity, danger). Threat of wind damage has restricted silviculture and led to the use of truncated rotations with precautionary felling in anticipation of windthrow.

Forest managers therefore require guidance on the timing and magnitude of wind damage. A number of risk-minimising strategies can be applied (see Quine *et al.*, 1995) but these may be costly and so are best carried out only where the risk is high. The strategies can involve operations within individual stands (typically 1–20 ha), or the layout of stands at the forest scale (areas up to 50 000 ha). A sound choice between techniques which may influence the risk of damage such as type of cultivation, thinning, sequence of felling, is most likely where prediction of their effects is possible. Research into both prevention and prediction of damage has been carried out for nearly 30 years and this past work has been drawn together to provide ForestGALES as a tool to guide forest managers.

The degree of constraint posed by strong winds in Britain

Wind damage is a serious problem in forests planted on exposed sites in Britain and western Europe. It is believed to cost the countries of the EU more than €15 million per year, and in extreme cases substantially more. Five ‘catastrophic’ storms have affected British forests since 1945, emphasising the frequency and scale of the problem – these include the storms of 1998 in south Scotland, 1990 in south-west England and south Wales, 1987 in the south-east of England, 1968 in the central belt of Scotland, and 1953 in north-east Scotland. In each case over 1 million cubic metres of timber was damaged, and up to five times the normal annual cut for the affected region required harvesting. In 1999, storms damaged more than 190 million cubic meters of timber in continental Europe and in 2005 storms damaged more than 80 million cubic metres of timber across Northern Europe. Lesser storms also cause some wind damage in most years, and their combined effect is a serious constraint in upland forests. Windthrow has been the main form of damage, but wind snap can be locally important when trees are particularly well anchored or the crowns are loaded with wet snow.

Historical context – previous predictive windthrow model

Until recently choices of windthrow-minimising strategies were guided by the windthrow hazard classification ((WHC): Miller, 1985) which provided a method to zone forest areas of 500 ha or more by adding scores for windiness and soil together to estimate a hazard class. Each class was associated with a critical and a terminal height (see Table 1.1) which respectively indicated the heights at which damage was expected to start and to reach a level necessitating clearance. The WHC provided a successful basis for comparing sites and guiding decisions on thinning, but did not predict the timing or amount of damage with acceptable accuracy (Quine and Bell, 1998). It was based on

subjective weighting of the influence of strength of wind and inadequacy of rooting on the timing of wind damage. While this synthesised understanding in the 1970s it could not be readily updated to incorporate new knowledge.

Research since the WHC was devised has improved understanding of root anchorage (Coutts, 1983; Coutts, 1986; Nicoll *et al* 2006), turbulence (Gardiner *et al.*, 1997), adaptive growth (Nicoll and Ray, 1996), and wind climate (Suárez *et al.*, 1999). While it was possible to incorporate such advances in qualitative advice (Quine *et al.*, 1995) a new system was needed to treat wind risk quantitatively.

Table 1.1

Critical and Terminal heights for each of the six classes defined by the Wind Hazard Classification.

WHC	Critical height (m)			Terminal height (m)		
class	Non-thin	Selective	Line thin	Non-thin	Selective	Line thin
1	Unconstrained by windthrow			Unconstrained by windthrow		
2	25.0	22.0	21.0	31.0	28.0	28.0
3	22.0	19.0	17.0	27.0	25.0	23.5
4	19.0	16.0	14.0	24.0	21.5	18.5
5	16.0	13.0	12.0	19.5	17.5	15.5
6	13.0	10.0	9.0	15.5	13.5	11.5

What does ForestGALES calculate?

ForestGALES calculates the probability of *average trees* being damaged within a stand. Damage to the average tree will by implication mean that the stand as a whole will be substantially damaged.

How does ForestGALES compare to the WHC?

ForestGALES estimates the chance (or probability) of windthrow or stem breakage, rather than stating a precise height at which damage will occur as in the WHC. Probabilistic predictions are more realistic than precise heights since the occurrence of damaging winds varies from year to year, which has a powerful influence on the occurrence and spread of damage.

The risk of damage is extremely dependent on the windiness of the site. In the WHC the measure of windiness is much coarser than is used in ForestGALES. This allows ForestGALES to discriminate several levels of risk for trees in similar WHC classes.

For example, a crown thinned stand of YC 12 Sitka spruce planted on a deep ploughed gley soil, with average drainage, where the DAMS score is 14, will have a critical height of 19.0 m and a terminal height of 21.5 m. The standard Forestry Commission yield models (Edwards and Christie, 1981) indicate that the trees will reach 21.4 m top height at age 51 when the return period for damage will be 7 years. On the other hand if the DAMS score is 10, then the site will still have a WHC class of 4, yet the return period for damage to a 51-year-old stand has increased to 200 years.

What is new in ForestGALES 2

ForestGALES 2 represents a complete rewrite of the ForestGALES software. The new software is a much more efficient and robust code, which is easier to understand and to document. Furthermore, it allows much easier integration with other modelling tools. The main part of ForestGALES (the part that does the actual calculations) has been constructed as a stand-alone dynamic link library (DLL). This means that all the functionality of ForestGALES is available from any other programme. The creation of a DLL has made it much easier for ForestGALES to be integrated with GIS software such as ArcView. Currently trials are under way in a forest district on the use of ForestGALES within ArcView. Following these trials a general release of a ForestGALES add-on to ArcView will be made available.

Wind Climate

The most important change to the calculation of risk is the change to the wind climate calculator. The calculation of the probability of a particular wind speed occurring at a site is based on the DAMS score. Additional analysis of wind climate data from Forest Research wind monitoring areas has suggested that the wind climate calculator was pessimistic. The differences are most pronounced at low DAMS scores but even at a DAMS value of 15 the new regression has the effect of reducing the DAMS score by 2 to an effective score of 13.

This is the only change in the calculation of risk. Critical wind speeds remain identical to ForestGALES 1.3 but the probability of these wind speeds occurring is reduced.

Appearance

The appearance of ForestGALES has been improved to provide a more consistent view and to provide for easier operation.

Defaults

There is now a Defaults button (Alt + D) that can be used to store the current values as the default. Therefore, if the current soil is a brown earth and the Defaults button is pressed then brown earth becomes the default soil used in ForestGALES. This is particularly useful if one is consistently working with a particular combination of species, cultivation and soil types.

Multiple Stands (Batch) Mode

The Batch operations have been substantially improved. Entering data in Multiple Stands Mode now uses the same forms as for Single Stand data entry and there are three versions of Multiple Stands Mode matching the three methods of calculating risk (Predictions using field measurements, Predictions using yield models, Predictions through time). Unlike the previous version of ForestGALES it is now possible to use the yield models to enter data or to calculate risk through time when in Multiple Stands Mode. In all cases input data and results can be stored in files for later use.

Export

There are improved export facilities in ForestGALES 2.1 allowing results to be sent to Word® as a Doc or Rtf file in the Single Stand mode and to Excel® in Multiple Stands Mode for display, storage and printing.

System requirements

ForestGALES will run on IBM compatible personal computers that have a minimum of:

- Microsoft Windows operating system
- Pentium 133 megahertz (MHz) or higher processor
- 32 Mb of RAM
- CD-ROM drive
- 20 Mb of hard disk space if DAMS scores left on CD (520 Mb for full installation)
- Super VGA (800 x 600) display or higher-resolution monitor with 256 colours

Recommended: Microsoft Excel/Word 97® or above to use data export facilities

Installing ForestGALES

To install ForestGALES:

1. Place the ForestGALES CD in the CD drive.
2. The program will automatically install.

If automatic installation does not occur, then:

3. Press the **Start** button.
4. Press **Run**.
5. Type **D:\Setup.exe** into the dialog box, which will appear (replace the letter D with whatever drive letter corresponds to your CD-ROM drive).
6. Left click on **OK**.

By default the program will then be installed into the directory **C:\Program Files\Forest Research\ForestGALES_ver2_1** on the drive, although it may be placed elsewhere if required. The installation program will automatically place all the files where they are needed, and add an option to run ForestGALES from the Programs bar.

Uninstalling ForestGALES

To uninstall the program:

1. Press **Start**.
2. Choose **Settings**.
3. Choose **Control Panel**.
4. Choose **Add/Remove Programs** then select **ForestGALES 2.1** from the list of programs.
5. Left click on **Add/Remove**.

This will remove all the components and data files of ForestGALES 2.1, with the exception of files created while using the program. Exceptions include saved/exported data files and new yield models.

Running the model for the first time

Start the program from the Windows **Start**. After the initial welcome screen you will be presented with the main menu window as shown in Figure 2.1. Choose **Mode, Single stand, Prediction using field measurements** and a new query form will appear as shown in Figure 2.2.

Set:

- Species to 'Sitka spruce'
- Cultivation to 'notched planting'
- Drainage to 'Average'
- Soil to 'Peaty Gley'
- Current spacing to 2.6 m
- Top Height to 18 m
- DBH to 19 cm
- DAMS score of 15

Press the **Run** button and the Return Period, Wind Damage Risk Status and Critical Wind Speeds for overturning and breakage will appear in the previously empty boxes at the bottom of the form. Return periods will be 200 years for both overturning and breakage. Change the top height to 19 m using the up arrow next to the top height edit box and press the **Run** button again. The return periods will change to 53 years for breakage and 181 years for overturning. **You are running ForestGALES!**

The various options available for the model are explained in later sections of this manual.

Figure 2.1 The ForestGALES main menu.





Figure 2.2 Query form for Single stand predictions using field measurements.



How does the model work?

3

The model uses data relating to individual trees to estimate the risk of damage to stands of trees by answering three questions:

- 1. What force would be needed to uproot or break the tree?*
- 2. What wind speed would create the force required to damage the tree (i.e. what is the threshold or critical wind speed)?*
- 3. What is the probability of the threshold wind speed being exceeded?*

What force would be needed to uproot or break the tree?

The model calculates the strength of the stem and the resistance of the tree to overturning independently.

Stem strength is based on theoretical work regarding tree stems as structures, and a knowledge of the wood strength and stiffness of different tree species.

Resistance to overturning is based on an extensive database of tree pulling experiments that relate tree characteristics to the load required to pull a tree over. These experiments have been conducted on a range of soil and cultivation types for many species. However in some circumstances no data are available, so assumptions have been made to allow data to be extrapolated. When this happens, a warning message box will appear and the results should be used with caution. The effect of drainage intensity is included in the calculation based on the average change in force needed to pull over trees on well or poorly drained soils in comparison to soils with 'average' drainage.

What wind speed would create the force required to damage the tree?

In ForestGALES the wind loading on trees is calculated from the drag the forest exerts on the flow above the canopy. The drag is a function of the wind speed and the aerodynamic roughness of the canopy which is dependent on the crown size and the area of ground occupied by each tree.

Crown size is modelled using regressions based on the height and diameter of the tree.

From a knowledge of the average wind loading on each tree and the resistance to breakage and overturning, the wind speed that would just cause the tree to overturn or break is calculated. This includes calculating the extra force due to the overhanging weight of the crown and stem as the tree bends.

What is the probability of a damaging wind speed occurring?

Having calculated the wind speed required to damage the tree, the probability of such a wind speed is estimated.

The average windiness of the site is measured using the DAMS system, which is based on location, elevation and exposure. However, average winds are unlikely to damage the tree, and the risk of extreme winds that are sufficiently strong to cause damage must be derived from DAMS using a Weibull distribution.

This distribution is extremely sensitive, and small changes in the wind speed required to cause damage can be associated with large changes in the probability of damage occurring.

Having calculated the probability of damage, this is converted to a return period for a damaging wind speed occurring. This is the average interval between storms that are associated with damaging winds.

Wind Damage Risk Status provides a rough estimate of risk, ranging from Status 1 (return period for damaging winds more than 100 years) to Status 6 (return period less than 10 years).

Running ForestGALES

There are several ways to start the program.

The easiest is to press Start, followed by Programs, ForestGALES 2.1, ForestGALES. When this is done, the program will start and the main menu (Figure 2.1) will load onto the screen.

ForestGALES has been designed as a complete Windows application. Menus are easily accessible with the mouse in the top part of the main window, or they can be accessed using pop-up windows, shortcut keys or speed buttons.

Operating modes

ForestGALES can be used interactively in two operating modes:

- | | |
|----------------------------|--|
| Single stand mode | - calculates the risk for one particular stand. |
| Multiple stand mode | - calculates the risk for a number of stands one after another (batch mode). |

There are three ways of making predictions within these modes:

- | | |
|---|---|
| Predictions using field measurements | - calculates the risk of damage at a single point in time from stand characteristics defined by the user. |
| Predictions using yield models | - calculates the risk of damage at a single point in time from stand characteristics contained in yield models. |
| Predictions through time | - calculates the risk of damage over a typical rotation from stand characteristics contained in yield models. |

Selecting a mode and opening a query form

Having started the program, to select a particular mode, left click on Mode at the top of the main menu.

A menu will appear. Left click on the mode that you want, and the menu will disappear, and a form will appear automatically.

Alternatively, a new form can be created using the shortcut buttons at the top of the main window. The meaning of each shortcut button is indicated by a hint message that appears when the mouse is positioned over the button.

It is possible to have several copies of a query form open at the same time. This may be useful if you wish to compare alternative scenarios.


ForestGALES query forms

ForestGALES collects data from and writes results to query forms. The exact layout of the form depends on the mode (single stand, multiple stands) in which the model is being used and the type of prediction being made (using

Using ForestGALES

field measurements, using yield models, through time). Figure 4.1 shows the layout of the form for *Single stand predictions using field measurements*.

Figure 4.1. Query form for Single stand predictions using field measurements. The parts of this form are:



Stand characteristics box – describes the soil, spacing, cultivation, drainage, and (where appropriate) planting year.

Tree characteristics box – describes the size of the trees within the stand being modelled.

Upwind edge effect box – states whether a new brown edge is present, and, if so, the size of the gap created.

DAMS box – describes how windy the site is. DAMS can be entered directly, calculated exactly, extracted from a pre-calculated data file for a particular location or estimated roughly. The layout of the box depends on which selection is chosen.

Controls box – allows the user to run the model, save inputs, load a saved file, print results, close the form and obtain help.

Wind damage risk box – indicates the risk of uprooting or stem breakage occurring, and the estimated return period.

The query forms for other modes are described in later sections of this manual.

Navigating ForestGALES

Using the mouse to make selections

As the mouse is moved over the query form, any of the components can be selected by clicking the left mouse button. If text boxes (such as Stand ID), up/down boxes (such as spacing) or menu boxes (such as species) are selected, then changes can be made to the box by either typing in text (for the text and up/down boxes) or by selecting the arrow buttons at the right edge of the box (for menus and up/down boxes).

When one of these boxes is selected then pressing F1 will result in Help being displayed to describe the use of the box.

If the mouse is moved over one of the buttons then the appearance of the button will change. Clicking the left hand mouse button will cause the button on the form to be pressed. This can be used to Run the model, Print the form, etc.

Using the Tab key

Pressing Tab causes the 'focus' to move through each of the controls in turn. The appearance of focus depends on the type of control. For text boxes, up/down boxes and menus the focus is indicated by the background becoming blue. For the buttons, focus is indicated by a thin black border and a black dashed box around the caption (and picture, if present) on the button.

When a control has focus, pressing F1 will result in help being displayed that relates to the control.

A control which has focus can also be edited, changed or activated as follows:

Text boxes (e.g. Stand ID)	Edit using the keyboard;
Up/down boxes (e.g. spacing)	Edit using the keyboard;
Menu boxes (e.g. species)	Edit using the up & down keys;
Radio buttons (e.g. windfirm/brown edge)	Select using the up & down keys;
Buttons (e.g. Calculate DAMS or RUN)	Use the Enter key to press the button.

Using shortcuts

Shortcuts allow a combination of the ALT key plus a letter to be used to select an option.

The Alt + key shortcuts in the main menu are:

Alt + F	Open File menu
Alt + M	Open Mode menu
Alt + A	Open DAMS menu
Alt + O	Open Options menu
Alt + W	Open Window menu
Alt + H	Open Help menu

Within ForestGALES, the shortcut for a particular button is indicated by an underlined letter in the button caption.

Getting help

Help can be obtained either by clicking on a Help button, selecting Help from the main menu, or by pressing F1.

Pressing Help starts the help system – this allows the user to search for a particular topic or keyword.

Pressing F1 will display help relating to the component that has focus (usually the last item where you clicked the mouse). These help pages are therefore 'context sensitive'.

Introduction

This is the standard type of prediction. It allows the user to calculate the risk of damage based on the stand's mensurational characteristics at a **single** point in time.


The user enters data that describe the site, the tree crop and whether or not a new edge is present.

In individual stand predictions, pressing **RUN** will then calculate the probabilities of damage occurring based on the selected options. Making predictions for multiple stands is explained in Chapter 8.

The stand characteristics box

The stand characteristics box (Figure 5.1) allows you to describe the stand for which you wish to calculate the risk of damage. The options are described below.

Figure 5.1 The Stand characteristics box.



Stand Characteristics

Stand ID: ForestGALES

Cultivation: Notched Planting

Drainage: Average

Soil type: Peaty Gley

Current Spacing (m): 2.6

Current Stocking (N/ha)

Options – Stand ID

An identification for the stand. Note: names longer than 25 characters will not print correctly if a report is produced.

Options – Cultivation

The cultivation type of the stand. The options are grouped into three main categories, which are shown in Table 5.1.

Table 5.1

Cultivation methods available within ForestGALES, indicating the pooling of options used by the model. (Notched planting means no cultivation.)

Mounding	Shallow ploughing	Deep ploughing
Notched planting	Shallow ploughing <45cm	Deep ploughing >45cm
Turf planting	Scarifying	Disc trenching
Moling		Alternate single/double
Mound planting		Contour ploughing
Complete ploughing		

Predictions using field measurements

Options – Drainage

This describes how well drained the site is. Poor drainage (due to blocked drains), for example, results in increased risk of uprooting. The options are:

Average – site is ‘typical’

Poor drainage – site is much wetter than you would expect

Good drainage – site is much drier than you would expect

Options – Soil type

This describes the main soil type in the stand. The most important factor is whether the soil is a peat, a gley, or a freely draining soil. The options are shown in Table 5.2.

Table 5.2

Soil types available within ForestGALES, indicating the pooling of options used by the model. ‘Other’ soils are currently treated like gleys.

Free draining	Peats	Gleys	Other
Brown earth	Juncus (Flushed basin) bogs	Ground-water gley soils	Rankers and skeletal soils
Podzols	Molina (Flushed blanket) bogs	Peaty gley soils	Littoral soils
Ironpan soils	Sphagnum (flat raised) bogs	Surface-water gley soils	Man-made soils
Calcareous soils	Unflushed blanket bogs		
	Eroded bogs		

Options – Current spacing

The average spacing between trees *at the time of risk assessment*. It must be between 0.6 and 10 metres.


Options – Current stocking

The number of trees per hectare *at the time of risk assessment*. The user has the choice between this option and the previous one.

The tree characteristics box

The characteristics of the average tree of the stand are entered in the *Tree characteristics box* (Figure 5.2).

Figure 5.2 The Tree characteristics box.



The screenshot shows a software window titled "Tree Characteristics". It contains three input fields: a dropdown menu for "Species" set to "Sitka Spruce", a spinner box for "Top height of stand (m)" set to "18.0", and another spinner box for "Mean DBH (cm)" set to "19.0".

The options that can be selected from the Tree characteristics box are:

Options – Species

The main species in the stand. Options are:

Scots pine	Douglas fir
Corsican pine	Noble fir
Lodgepole pine	Grand fir
European larch	Sitka spruce
Japanese larch	Norway spruce
Hybrid larch	Western hemlock

Options – Top height

Top height (in metres) of the stand being assessed. This must be between 5 and 75 m.

Options – DBH

Mean diameter (in cm) of the stand being assessed. This must be between 5 and 50 cm.

Predictions using field measurements


The DAMS box

DAMS is a measure of site windiness. The greater the value, the windier the site. Only values between 5 and 32 (the range of values found in Britain) will be accepted by the model. If the DAMS score for the site is known then it can be entered directly. If it is not known then it can be estimated, looked up or calculated by the program. The DAMS box is displayed in Figure 5.3.

Grid References method

In the DAMS box, the user can select **Grid Reference** and then enter the grid reference of the site; the DAMS score, if available, will be displayed. The DAMS scores are supplied on a separate CD-ROM, and this must be installed in the CD drive if the data are to be accessed.


Figure 5.3 The DAMS box.



Exact Calculation method

In the DAMS box, select **Calculation** then click the **Apply...** button. The calculation box will appear. Selecting the **Exact Calculation** box (Figure 5.4) will give the opportunity to calculate the exact DAMS score. The Wind Zone for the location can be found using the map that is displayed when **View Map** is pressed. Elevation is entered together with the Topex values for each of eight compass directions (TOPEX is the angle to the horizon in whole degrees in the particular compass direction, with values less than zero being entered as zero). The resulting DAMS score can be copied to the main form by pressing **Apply**, or discarded by pressing **Cancel**.


Figure 5.4 The Exact Calculation box.



Rough guess method

In the DAMS box, select **Calculation** then click the **Apply...** button. The calculation box will appear. Selecting the **Rough Guess** box (Figure 5.5) will give the opportunity to estimate the DAMS score. Select the options that best describe the site based on region within GB, elevation, shelter and aspect. Press **Apply** to copy the resulting DAMS score to the query form, or **Cancel** to close the window without copying the value across. This method is particularly useful for making general comparisons between sites.

Figure 5.5 The Rough Guess box.




The image shows a software dialog box titled "DAMS Calculation". It has two tabs: "Rough Guess" (which is selected) and "Exact Calculation". Under the "Rough Guess" tab, there are four dropdown menus: "GB Region" (set to "West & North Coast Scotland & Islands"), "Elevation" (set to "Top of Hill"), "Shelter" (set to "Well Sheltered"), and "Aspect" (set to "All"). To the right of these menus, there is a "Result" section with a label "DAMS" and a text box containing the number "23". Below the "Result" section is a "Controls" section containing three buttons: "Apply", "Help", and "Cancel".

The upwind edge effect box

The *Upwind Edge Effect* box (Figure 5.6) is used to describe whether a new non-wind-firm edge has been created adjacent to the stand being modelled. Brown edges (i.e. edges that were originally not at the stand edge) are often a place where wind damage starts. If a new edge has been created then the **Brown Edge** button should be pressed. The size of the gap can then be altered. The default value is 0 m. The effect of a gap increases with gap width until the size equals 10 x mean tree height, after which the effect remains at a maximum.

Figure 5.6 The Upwind Edge Effect box.




The image shows a software dialog box titled "Upwind Edge Effect". It has two radio buttons: "Windfirm edge" (which is selected) and "Brown edge - Size of gap (m)". The "Brown edge" option has a small text box next to it containing the number "0".

Predictions using field measurements

The controls box

The *Controls* box (Figure 5.7) allows the user to control ForestGALES. The options are described below.

Figure 5.7 The Controls box.



The wind damage risk box

Model outputs are displayed in this box (Figure 5.8).

Figure 5.8 The Wind Damage Risk box.



Critical wind speed

The critical wind speeds (i.e. the lowest wind speeds that will cause overturning or breakage) are presented. By default these are presented in mph. Wind speed units can be changed in the **Options** menu.

Return period

The *average length* in years between wind speeds exceeding the critical wind speed occurring at the site. This is the *average* interval between gales that will damage the site. The statistical nature of the wind climate means that strong winds with long return periods (i.e. 50 years) may occur within a few years of each other. The occurrence of a damaging storm does not alter the risk of further damage in subsequent years.

Wind damage risk status

A measure of the risk of damage either by overturning or stem breakage. Six classes of wind damage risk status exist. These are listed in Table 5.3.

Table 5.3

Wind risk status and associated return periods.

	Return period
1	>100 years
2	100–50 years
3	50–33 years
4	33–20 years
5	20–10 years
6	<10 years

Unlike the Windthrow Hazard Class (WHC) classes, the risk status of a site will change over time. If the risk status for stem breakage is greater than for overturning, then stem breakage is likely to be the predominant form of damage, whereas if the risk status is greater for overturning than for breakage then uprooting is likely to be the predominant form of damage.

Windthrow Hazard Classification (WHC)

The WHC class for the site, based on the 'traditional' system (Miller, 1985) using the DAMS score and soil type. This is provided to allow comparisons to be made between the old and new system.

Predictions using field measurements

Introduction

This type of prediction allows the user to calculate the risk of damage at a single point in time based on stand characteristics defined in yield models. It is particularly useful when the mensurational characteristics of a stand are not known or when wanting predictions made at a single point in the future. Figure 6.1 shows the *Predictions using yield models* query form for single stands. Making predictions for multiple stands is explained in Chapter 8.

A selection of yield tables is distributed with ForestGALES, and is stored in the directory `\yldmdl`s. To select a yield model, choose the species, then select the yield class, thinning regime and initial stocking that apply. Only the yield models which are available can be selected from the program. It is, however, possible to create new yield models which can be used as input for ForestGALES. This is described on page 6.3.

Figure 6.1 The query form for Single stand predictions using yield models.

Stand characteristics box

As for *Predictions using field measurements*, the *Stand characteristics* box contains the **Stand ID**, **Cultivation**, **Drainage** and **Soil type** options (see page 5.1). The current stocking cannot be specified as this is taken from the yield tables.

Options – Planting year

A planting year can be specified. This will automatically adjust the stand age to present in the *Tree characteristics* box. If you want to make a prediction into the future adjust the age of the stand (see below) instead of specifying a planting year.

Tree characteristics box

When predictions are made using yield models, the *Tree characteristics* box shows the same **Species** option as found in the *Predictions using field measurements* (see page 5.3). However, the other options are different:

Options – Yield class

The thinning model regimes displayed will depend on the yield models available (Edwards and Christie, 1981).

Options – Thinning regime

- Intermediate thinning with no delay
- Intermediate thinning with five year delay
- Intermediate thinning with ten year delay
- Line thinning
- Line thinning with five year delay
- Line thinning with ten year delay
- Crown thinning
- No thinning

Options – Initial spacing

The initial spacing (in metres, based on square planting) should be selected. The range varies from 0.9–3.0 m depending on yield models available.

Options – Age

The age of the crop to be modelled. The valid range varies between yield models. Ages between published values are calculated using linear interpolation between the nearest younger and nearest older published value. By default if the date of planting has been entered then the age of the stand will be calculated. This will give the risk now. It is possible however to change the date, allowing the risk to be calculated at some other time.

The 'Tree Details' button

If this button is pressed then the height, diameter, current spacing and volume of the trees will be displayed. This is designed to provide information about the type of stand being modelled.

Other boxes

The **DAMS** box, the **Controls** box, the **Upwind edge effect** box and the **Wind damage risk** box are identical to those described on pages 5.4 to 5.7.

User-defined yield models

New yield models can be constructed for use within ForestGALES using a word processor, or spreadsheet program.

Overview

Currently ForestGALES doesn't contain a user-defined yield model helper. Yield models must therefore be created using a word processor or Excel®. The general layout of the yield model file is described below. An Excel® template is included with ForestGALES in directory `yldmodls\userdefined` and is called `yieldmodel.xls`.

The format of the model is illustrated in Table 6.1 and is:

Line one: a header to describe what is in each field

Line two onwards: fields separated by a single space to define

- Age (years)
- Top Height (m)
- Stocking density (stems ha⁻¹)
- DBH (cm)
- Basal area (m² ha⁻¹)
- Mean tree volume (m³/tree)
- Volume per hectare (m³ ha⁻¹)

In each case data refer to the main crop after thinning. This is the format of the Forestry Commission Yield Models.

Table 6.1

Layout of a yield model for use in ForestGALES.

Age (years)	Top height (m)	Trees/ha	Mean DBH (cm)	Basal area (m ² /ha)	Mean tree volume (m ³)	Volume (m ³ /ha)
20	7.4	2781	11	26	0.03	71
25	9.2	2300	13	32	0.06	90
30	10.9	1900	15	38	0.10	120

Naming user-defined yield models

The file should be saved as a text file with a file extension of `.yld`.

If ForestGALES is to recognise the model then it must be named in a specific way. This consists of an 8 character name.

1. The first two characters indicate species; these are shown in Table 6.2.
2. The second two characters refer to the yield class (02–30).
3. The third two characters refer to thinning regime, and are shown in Table 6.2.

Predictions using yield models

- The final two characters refer to initial spacing in metres x 10. Therefore 0.9 m spacing becomes 09, and 2.1 m spacing becomes 21.

Table 6.2

Species and thinning codes for naming user-defined yield models.

Species code	Species	Thinning code	Thinning regime
SS	Sitka spruce	IZ	Intermediate thinning no delay
NS	Norway spruce	IF	Intermediate thinning five years delay
SP	Scots pine	IT	Intermediate thinning ten years delay
LP	Lodgepole pine	LZ	line thinning no delay
CP	Corsican pine	LF	line thinning five years delay
EL	European larch	LT	line thinning ten years delay
JL	Japanese larch	CZ	crown thinning
HL	Hybrid larch	NO	non-thinning
DF	Douglas fir	T1	user-defined thinning regime
GF	Grand fir	T2	user-defined thinning regime
NF	Noble fir	T3	user-defined thinning regime
WH	Western hemlock		

A user-defined model for yield class 18 Sitka spruce for a non-standard thinning regime planted initially at 2.0 m spacing would therefore be saved as **SS18T120.yld**.

The file should be saved in the directory **yldmdls\XX** where **XX** is the two letter species code indicated in Table 6.2.

If a new model is created with an identical name to a model that already exists, then the old model will be lost.

Using Excel® to create user-defined yield models

- Open the file **yieldmodel.xls** in Excel®. It is in the directory **\yldmdls\userdefined** wherever you have installed ForestGALES.
- Type data for the yield model into the template.
- Extra lines can be added as necessary. If the template contains more lines than are needed, then remember to delete the extra lines.
- Press **File, Save As**.
- Choose the option **Formatted Text(Space delimited)(*.prn)**.
- The file must be saved in the directory **XX**, where **XX** is the two character species code indicated in Table 6.2.
- Type the filename according to the format described on page 6.3 to 6.4.

Predictions using yield models

Note: The filename must be enclosed in quotes (e.g. "SS20IZ20.yld"), otherwise the file extension .prn will be added resulting in a filename called something like SS20IZ20.yld.prn

8. Press **Save**.

The new model will be accessible within ForestGALES when a new *Predictions using yield models* query form is opened.

Introduction

This type of prediction allows the user to calculate the risk of damage to a stand over part or all of its rotation, rather than at just a single age.

Figure 7.1 shows the *Predictions through time* query form for single stands. Making predictions for multiple stands is explained in Chapter 8. The main difference between predictions through time and predictions at a single point in time is that the results are displayed on a separate form when the model is run. Also, there is no option for modelling the effect of a new edge.

Figure 7.1 The query form for single stand predictions through time.

Stand characteristics box

This is identical to the Stand characteristics box in Predictions using yield models (see page 6.1) except that year of planting is not available, since Predictions through time calculates risk over the whole rotation, not at just a single age.

Tree characteristics box

This is identical to the Tree characteristics box described on page 6.2 except that age is not available.

The DAMS box

This is identical to the DAMS box described on page 5.4.

The controls box

This is identical to the Control box described on page 5.6, except that a report cannot be created.


User-defined yield tables

New yield tables can be constructed using the method described on page 6.3.

Viewing the results

Figure 7.2 shows a typical results form. The form has four areas: a graphics window, two tabular windows and a controls box. These are described below.

Figure 7.2 The Results form for single stand predictions through time.



The graphics window

The graphics window indicates the return period in years for damaging storms at intervals throughout the rotation. Typically as the trees grow older and taller the risk of wind damage increases and the return period therefore decreases.

The graph contains two lines. The red line indicates the risk of overturning, and the blue line indicates the risk of stem breakage. The data relate to average trees (i.e. with mean diameter and height) based on the yield tables.

Details of the graph can be investigated as follows:

Zooming into the graph:

- Place the mouse cursor at the top left hand corner of the area you wish to zoom in to.
- Hold down the left mouse button.
- With the mouse button held down, move the mouse to the lower right hand corner of the area you wish to zoom in to.
- Release the mouse button.
- The display will then change to the selected area.

Note: there is a minimum area which can be selected, and if a smaller area is selected then no zooming will occur.

Scrolling around the graph:

- Place the mouse cursor somewhere on the graph.
- Hold down the right mouse button.
- With the right button held down, move the mouse.
- The graph will move as you do this.

Resetting the graph:

- Place the mouse cursor on the graph.
- Hold down the left hand mouse button.
- With the left button held down move the mouse to the left and upwards.
- The display will then revert to the original state.

Note: there is a minimum area which can be selected, and if a smaller area is selected, then the graph will not reset.

Displaying new results:

The effect of changing the yield model used can be observed directly in the graphics window. Go to the query form, change any parameter, press the Run button and the graphics display will change. This allows the user to easily observe the effect of changing the thinning model or soil type or species, for example.

The tabular data windows

The table window on the left hand side indicates how return period changes with age, and provides details on height, diameter, spacing and volume of the crop with age. The date when the crop enters each Wind Damage Risk Status (WDRS) is displayed in the table window on the right hand side. Data can be exported as described on page below.

The results form controls box

The controls that can be used on the results form are:

Print Form	Prints the current query form. Shortcut is Alt + P .
Report	Creates a report that contains all the information on the form. The report can be printed or saved for later use. Shortcut is Alt + T .
Help	Starts the Help system. Shortcut is Alt + H .
Excel® Export	Allows the user to export the tabular results to Excel®. Shortcut is Alt + X .


Introduction

ForestGALES has been designed to allow batch files of data (Multiple stands) to be processed, in addition to assessing the risk to stands one at a time. This allows data for a whole forest, property or coupe to be stored and calculated together.

In this mode, data are read in from one file, and output to a different file. This can be very helpful if a large number of stands need to be assessed at a single time. Those stands would not necessarily have to be linked to each other as the calculations are made independently. The *Multiple stands mode* can be used with each type of prediction.

The interface of ForestGALES in *Multiple stands mode* is different from that used for individual stands, and contains input and output areas as well as the control panel. Figures 8.1–8.3 show the interface for each type of prediction.

Figure 8.1 The query form for Multiple stand predictions using field measurements.



Multiple stands

Figure 8.2 The query form for Multiple stand predictions using yield models.





Figure 8.3 The query form for Multiple stand predictions through time.



Input files can be created using the *Multiple stands helper* window that appears in front of the query form when working in the *Multiple stands mode* (see Figure 8.1–8.3). The input files can also be created from a spreadsheet. However, since special codes are needed for some parameters, the use of the *Multiple stands helper* application is recommended.

The multiple stands helper

The multiple stands helper allows the user to select inputs from menus, and these are then written to the input area of the query form. This is useful because ForestGALES uses coded values to describe the species, site and cultivation rather than descriptions, and these values may be difficult to remember. For each prediction type, the options are identical* to those described for the corresponding query form in the *Single stand* prediction mode (See Sections 5, 6 and 7).

*Note: Clicking on **Close** causes both the helper and the Multiple stands query form to be closed. Unless you want to discard the inputs, click on **Calculate Risk** before closing the helper.*

*There is a slight difference when making predictions for multiple stands using yield models. The year at which you wish to calculate the risk has to be specified on the query form rather than in the helper.

The controls box

The controls in the *Multiple stands helper* are:

Add	Adds the selected values to the input area of the query form. Shortcut is Alt + A .
Print Form	Prints the current query form. Shortcut is Alt + P .
Help	Starts the Help system. Shortcut is Alt + H .
Defaults	Changes the default values in the registry to the ones included in the query form. Shortcut is Alt + D .
Close	Closes both the helper and the query form. Shortcut is Alt + C .

The multiple stands query form

The characteristics of the stand are entered in the **input area**. All data relating to a single stand are displayed on the same line. After running the program, the results of the model are displayed in the **output area**. If the data have been saved then the filename will be displayed at the top of the output area. The outputs vary slightly depending on the mode you are running in (see Table 8.1).

Table 8.1

Modes available for multiple stand predictions

MODE	OUTPUTS
Predictions Using Field Measurements	Return Period for Overturning, Wind Damage Risk Status for Overturning, Return Period for Breakage, Wind Damage Risk Status for Breakage.
Predictions Using Yield Models	Model Used, Current Top Height (m), Current DBH (m), Current Spacing (m), Return Period for Overturning, Wind Damage Risk Status for Overturning, Return Period for Breakage, Wind Damage Risk Status for Breakage.
Predictions Through Time*	Model Used, Age to reach WDRS 4 for overturning, Age to reach WDRS 5 for overturning, Age to reach WDRS 6 for overturning, Age to reach WDRS 4 for breakage, Age to reach WDRS 5 for breakage, Age to reach WDRS 6 for breakage.

*Double clicking on any of the input lines in this mode will open the graphics display window as for a single stand. In this way it is possible to observe differences between the risk for all the stands entered as inputs.

The controls box

The controls in the *Multiple stands* query form are:

- Calculate Risks** Runs the model using the selected values. Shortcut is **Alt + R**.
- Delete Row** Deletes a row of data from the input area. Shortcut is **Alt + E**.
- Open File** Opens a previously saved query form. Shortcut is **Alt + O**.
- Save Inputs** Saves the input file for later use. Shortcut is **Alt + N**.
- Save Outputs** Save the output file so that it can be exported to another application. Shortcut is **Alt + U**.
- Help** Starts the Help system. Shortcut is **Alt + H**.
- Excel® Export** Allows the user to export the tabular results to Excel®. Shortcut is **Alt + X**.

Year of calculation

When you are in the *Predictions Using Yield Models mode* then the option is provided to modify the Year of Calculation in the bottom right corner. The default is the current year but any year in the past or future (up to 2100) can be chosen. If the age is less than the minimum in the yield table the model defaults to the minimum age. If the age is beyond the maximum age in the yield tables the model defaults to the maximum age.

9

Single stand predictions using field measurements – getting started

1. Start **ForestGALES**.
2. Left click on **Mode**
 - ⇒ The Mode dialog box will open.
3. Choose **Single stand, Predictions using field measurements**
 - ⇒ A new query form will open. This form has a range of pre-selected options. However, no value will be present in the Wind Damage Risk box.

*For the purpose of this exercise, make sure the default options are set to the following values: **Cultivation** Notched planting, **Drainage** Average, **Soil type** Peaty Gley, **Current spacing** 2.6 m, **Species** Sitka spruce, **Top height of stand** 18 m, **Mean DBH** 19 cm, **DAMS Score** 15. **Windfirm edge**.*
4. Left click on **Run**
 - ⇒ Values of 200 years will appear in the Wind Damage Risk box for the return period for both overturning and breaking.
5. Left click on the up button to the right of the **Mean DBH** box
 - ⇒ The DBH will increase by 0.1 cm per 'click'.
6. Keep changing the DBH in this way until a value of 25 cm appears in the box.
7. Change **Top height** until it reads 21 m.
8. Change **Current Spacing** until it reads 3.8 m.
9. Left click on **Run**
 - ⇒ The values in the Wind Damage Risk boxes will change to 12 years for **Return period for overturning** and 13 years for **Return period for breakage**.

This indicates that the risk of damage changes rapidly with changes in DBH, height and spacing. Other parameters can be changed in a similar way.

Single stand predictions using field measurements – the effect of a new edge

This example shows how creating new edges affects the risk of damage.

1. Start ForestGALES and open a query form for Single stands predictions

- using field measurements in the same way as the previous example.
- Using the pre-selected options, left click on **Run**
 - ⇒ This will give the return periods for damage for the default parameters, and no new edge.
 - Left click on the **Brown Edge** button
 - ⇒ A black dot will appear on the button, and the value for size of upwind gap will become black. Change the **Size of gap** to 400 m.
 - Using the pre-selected values left click on **Run**
 - ⇒ The values in the Probabilities boxes will change and should become 4 years for **Return period for overturning** and 3 years for **Return period for breakage**.

This indicates that the risk of damage is much greater if a new edge is present, than if no new edge is present. The Gap size box will also have changed to 167 m. This is 10 x the mean tree height. Gap widths greater than this have no additional effect on stability.

- Now try changing the size of the gap to 5 m and left click on **Run**
 - ⇒ The values in the Probabilities boxes will change and should become 13 years for **Return period for overturning** and 8 years for **Return period for breakage**.

Again, try changing other options to see what happens.

Single stand predictions using yield models

This example shows how to use yield models to provide input data for the model.

- Left click on **Mode**
 - ⇒ The Mode dialog box will open
- Choose **Single stand, Predictions using yield models**
 - ⇒ A new query form will open. This form has a range of pre-selected options. However, no value will be present in the Wind Damage Risk box.

*For the purpose of this exercise, make sure the default options are set to the following values: **Cultivation** Notched planting, **Drainage** Average, **Soil type** Peaty Gley, **Species** Sitka spruce, **Yield class** 8, **Thinning regime** Crown Thinning, **Initial spacing** 1.7 m, **Age** 37, **DAMS Score** 15. **Windfirm edge**.*

3. Left click on **Run**

- ⇒ This will give the probabilities of damage for a stand of YC 8 Sitka spruce, planted at 1.7 m spacing with a crown thinning regime at an age of 37. The values in the Probabilities boxes will change and should become 200 years for **Return period for overturning** and 200 years for **Return period for breakage**.

The size and spacing of the modelled trees can be viewed using the **Tree Details** button.

4. Now change the age to 55 and left click on **Run**

- ⇒ The values in the Probabilities boxes will change and should become 75 years for **Return period for overturning** and 81 years for **Return period for breakage**.

This indicates that the risk is increasing with stand age.

5. Now left click on the **down arrow** to the right of Thinning regime

- ⇒ A menu of thinning options will appear as you move the mouse over the options, they are highlighted in turn.

6. Left click on **Intermediate with no delay**

7. Left click on **Run**

- ⇒ The values in the Probabilities boxes will change and should become 200 years for the **Return period for overturning** and 200 years for the **Return period for breakage**.

Again, try changing other options to see what happens.

DAMS example

This example demonstrates the use of DAMS as an input for the model.

1. Start **ForestGALES** and open a query form for Single stands predictions using yield models in the same way as the previous example.
2. Leave the options as they were in the previous example and left click on **Run**

⇒ This will give the probabilities of damage for a typical tree in a stand of YC 8 Sitka spruce, planted at 1.7 m spacing with a crown thinning regime at an age of 37. The values in the Probabilities boxes will change and should become 200 years for **Return period for overturning** and 200 years for **Return period for breakage**.
3. Within the **DAMS** box, use the up and down arrows to select 19 as the

value of DAMS score. The higher the DAMS score the windier the site.

4. Leave the options as they are and left click on **Run**

⇒ The return periods should change to 23 years for the **Return period for overturning** and 57 years for **Return period for breakage**.

It should be apparent that relatively small changes in DAMS result in large changes in calculated risk.

Again, try changing other options to see what happens.

DAMS – rough guess example

This example demonstrates using a rough estimate of DAMS as an input for the model.

1. Start **ForestGALES** and open a query form for Single stands predictions using yield models.

2. Within the DAMS box select the **Calculation** button then click **Apply...**

⇒ A new form will appear.

3. Left click on the **Rough Guess** tab

⇒ New options will appear.

4. Select **GB Region** Scottish & English Borders, **Elevation** Mid-Slope, **Shelter** Exposed, **Aspect** West.

⇒ 18 will appear in the DAMS box Pressing **Apply** would copy this value to the query form.

5. In the **GB Region** box, select Central Wales

⇒ 17 will appear in the DAMS box.

Again, try changing other options to see what happens.

DAMS – grid reference example

This example demonstrates how to obtain the DAMS score for a specific grid reference.

1. Start **ForestGALES** and open a query form for Single stands predictions using yield models.

2. Within the DAMS box click the **Grid Reference** button. Press **Apply...**

⇒ 15 will appear in the DAMS box. This is the value of DAMS for

NH180150. If ForestGALES cannot find the DAMS data it will ask you to locate it. The **Find Files** option in the Windows start-up menu may be useful for this.

3. Type NY659932 in the Grid Reference box

4. Left click **Apply**

⇒ 17 will appear in the DAMS box.

Again, try changing other options to see what happens.

DAMS – calculate DAMS example

This example demonstrates how to obtain the DAMS score for a location based on ground measurements.

1. Start **ForestGALES** and open a query form for Single stands predictions using yield models

2. Within the DAMS box select the **Calculation** button then left click **Apply...**

⇒ A new form will appear.

3. Left click on the **Exact Calculation** tab

⇒ New options will appear. By default, 23 will appear in the DAMS box. This would relate to a hill top site at 200 m elevation in the west of Scotland. Pressing **Apply** would copy this value to the query form.

4. To change the location left click on **View Map**

⇒ This will cause a new window to appear.

5. Left click in the dark blue area in central Scotland (North West of the Tay estuary).

6. Press **OK**

⇒ The windzone value 3 should be placed into the **Windzone** box and 15 will appear in the DAMS box.

7. Select **100** in the Elevation box

⇒ 13 will appear in the DAMS box.

Table 9.1 Example Topex values.

8. Enter **Topex** scores as shown in Table 9.1.

North	0	South	5
North-east	3	South-west	2
East	5	West	0
South-east	10	North-west	0

11. Left click **Calculate**

⇒ 10 will appear in the DAMS box.

Again, try changing other options to see what happens.

Anchorage	The complex of mechanisms by which the root system and soil resist the wind forces on the stem and crown.
Brown edge	An edge of a stand that was created by felling part or all of the adjacent crop, rather than being a crop boundary since the time of planting.
Centre of pressure	The average position in the crown of the tree where the total force of the wind can be said to act.
Coherent gusts	Organised rotational motions in the air (= Vortices).
Critical wind speed	Threshold hourly wind speed above which the average tree of a stand is expected to be overturned or snapped.
Critical height	The top height of the stand at which damage was expected to start within the WHC system.
Cultivation	The method of preparing the soil prior to tree establishment. This may have been done by many means such as ploughing, mounding or producing turves.
Damping	The processes by which oscillations are reduced in size and tend to stop. Damping includes canopy clashing, canopy drag through the air, and frictional movement of stem fibres.
DAMS score	Detailed Aspect Method of Scoring – a system for scoring windiness derived from tatter flags and using representation of location and terrain to calculate a score (Quine and White 1993).
DBH	Diameter of a tree at 1.3 m above ground level.
Dominance	The 'social' status of a tree within a crop. Five categories are usually defined – dominant (trees with a crown entirely within the canopy), codominant (trees with much of their crown in the canopy), subdominant (trees with their crowns generally below the canopy), suppressed (trees with small crowns entirely beneath the canopy, which are gradually dying from lack of light) and dead.
Drag area	The surface area of the tree (canopy and stem) presented to the wind. Drag area is reduced as wind speed increases, due to streamlining of the tree.
Drag force	The force on the tree caused by the pressure exerted by the wind on the crown (= wind loading).
Drainage	A description of site wetness; poor refers to a wet site where rooting is severely restricted by a shallow watertable – due to local topography or failure of/lack of installed

	drainage system; good refers to better than average site – due to topography (e.g. shedding slope) or good quality intensive drainage network.
Frequency of oscillation	The number of sway cycles of the tree per second.
Fulcrum	The position on the lee-side of the tree where the root system pivots when the tree is bent by the wind (= hinge).
Gust	A rapid increase in wind speed over a short period of time (seconds rather than minutes).
Hinge	See Fulcrum.
Leeward	The side of the tree facing away from the wind.
Lever	The distance between the point of action of a force and the fulcrum.
Modulus of elasticity	A measure of stiffness. Modulus of elasticity is the load that theoretically would be required to make a material double (or halve) in length were it to behave perfectly elastically. In practice timber generally stops behaving elastically when its length changes by 1% and breaks when the change exceeds 2%.
Modulus of rupture	The force per unit area that is required to break a material when a bending load is applied.
Moment	Force multiplied by distance (= torque).
Overtipping moment	The force on the tree multiplied by the distance from where the force acts (the centre of pressure) to the fulcrum, plus the additional moment due to the weight of the over-hanging crown.
Risk (for a tree)	The probability in a particular year of the critical wind speed being exceeded (see Vulnerability).
Risk status	A measurement of the probability of the critical wind speed being exceeded in a particular year, grouped into six categories. Status 1: return period >100 years; Status 2: return period 50–100 years; Status 3: return period 33–50 years; Status 4: return period 20–33 years; Status 5: return period 10–20 years; Status 6: return period <10 years.
Root architecture	The appearance and structure of the root system, particularly the number and arrangement in three dimensions of the thickest roots.

Spacing – Current	The average spacing between trees at the time of risk assessment.
Spacing – Initial	The average spacing between trees at the time of planting.
Stocking	The number of trees per hectare at the time of risk assessment.
Terminal height	The top height of a stand at which wind damage was expected to reach a level necessitating clearance.
Thinning	The removal of a proportion of the tree crop for silvicultural or economic reasons. ForestGALES can extract data from yield models categorised according to thinning regime and initial spacing. The regimes used in the yield models are non-thinning (no trees actively removed, though some may die naturally), intermediate thinning (removal of trees from throughout the crop, with the smaller trees being preferentially removed), line thinning (removal of trees in straight lines) and crown thinning (removal of trees that are competing with the crowns of the highest quality trees). Models have also been produced for delayed thinnings.
Top height	The average height of the 100 trees of largest diameter per hectare, usually measured as the average height of the largest diameter trees in a sample of 0.01 ha plots.
Topex	A measure of exposure based on the sum of the angles to the horizon in eight compass directions.
Turbulence	The random variations in wind speed and direction.
Vortices	See Coherent gusts.
Vulnerability (of a tree)	The threshold wind speed required to blow over a particular tree on a particular site.
WHC	See Windthrow Hazard Classification.
Wind loading	See Drag force.
Windthrow hazard classification	A method to zone forest areas of 500ha or more by adding scores for windiness and soil together to estimate a hazard class. Each class was associated with a critical height and a terminal height.
Windward	The side of the tree facing towards the wind.
Wind zone	A range of windiness categories for the whole of Britain. The higher the wind zone the windier the climate on average. Windzone boundaries are defined in Quine and White (1993).

Glossary

Yield class

A commonly used expression of growth rate – defined as the mean maximum annual increment that could be achieved by the stand (units – $\text{m}^3 \text{ha}^{-1} \text{yr}^{-1}$).

General

Coutts, M. P. and Grace, J. (Eds). (1995).
Wind and trees.
Cambridge University Press, Cambridge, pp 485.

Peltola, H., Gardiner, B.A., Kellomäki, S., Kolström, T., Lässig, R., Moore, J.,
Quine, C. P. and Ruel, J-C. (2000).
Wind and other abiotic risks to forests.
Forest Ecology and Management **135**, Special Issue, pp 364.

Quine, C. P., Coutts, M., Gardiner, B. and Pyatt, G. (1995).
Forests and wind: Management to minimise damage.
Forestry Commission Bulletin 114.
HMSO, London.

Ruck, B., Kottmeier, C., Mattheck, C., Quine, C. and Wilhelm, G. (2003).
Wind effects on trees. Proceedings of the international conference.
University of Karlsruhe, Germany, September 16–18, 2003.

Wind

Barry, R. G. (1992).
Mountain weather and climate, 2nd Edition.
Routledge, London, pp 402.

Cook, N. J. (1985).
The designer's guide to wind loading of building structures.
Part 1: Background, damage survey, wind data and structural classification.
Butterworths, London, pp 371.

Finnigan, J. J. and Brunnet, Y. (1995).
Turbulent airflow in forest on flat and hilly terrain. In: *Wind and trees*, (Eds)
M. P. Coutts and J. Grace.
Cambridge University Press, Cambridge, 3–40.

Gardiner, B. A. (1994).
Wind and wind forces in a plantation spruce forest.
Boundary Layer Meteorology **67**, 161–186.

Kaimal, J. C. and Finnigan, J. J. (1994).
Atmospheric boundary layer flows; their structure and measurement.
Oxford University Press, pp 289.

Oke, T. R. (1985).
Boundary layer climates.
Methuen, London, pp 372.

Quine, C. P. and White, I. M. S. (1993).
Revised windiness scores for the Windthrow Hazard Classification.
Forestry Commission Research Information Note 230.
Forestry Commission, Edinburgh.

References & Further Reading

Stacey, G. R., Belcher, R. E., Wood, C. J. and Gardiner, B. A. (1994).
Wind and wind forces in a model spruce forest.
Boundary Layer Meteorology **69**, 311–334.

Suárez, J.C., Gardiner, B.A. and Quine, C.P. (1999)
A comparison of three methods for predicting wind speeds in complex
forest terrain.
Meteorological Applications **6**, 1–14.

Soils

Avery, B. W. (1990).
Soils of the British Isles.
CAB International, Wallingford. pp 463.

Kennedy, F. (2002).
The identification of soils for forest management.
Forestry Commission Field Guide.
Forestry Commission, Edinburgh pp 56.

King, J. A., Smith, K. A. and Pyatt, D. G. (1986).
Water and oxygen regimes under conifer plantations and native vegetation
on upland peaty gley soil and deep peat soils.
Journal of Soil Science **37**, 485–497.

Pyatt, D. G. (1987).
The effect of afforestation on the water and oxygen regimes of stagnopodzols.
North of England Soils Discussion Group Proceedings **22**, 93–108.

Pyatt, D. G. (1990).
Forest drainage.
Forestry Commission Research Information Note 196.
Forestry Commission, Edinburgh.

Pyatt, D. G., Anderson, A. R., Stannard, J. P. and White, I. M. S. (1985).
A drainage experiment on a peaty gley soil at Kershope Forest, Cumbria.
Soil Use and Management **1** (3), 89–94.

Pyatt, D. G. and Craven, M. M. (1979).
Soil changes under even-aged plantations. In: *The ecology of even-aged
plantations*, eds D. Ford, D. C. Malcolm and J. Atterson.
Proceedings of meeting of Division 1, IUFRO Sept 1978. ITE 1979, 369–386.

Pyatt, D. G. and Smith, K. A. (1983).
Water and oxygen regimes of four soil types at Newcastleton Forest, south
Scotland.
Journal of Soil Science **34**, 465–482.

Ray, D., White, I. M. S. and Pyatt, D. G. (1992).
The effect of ditches, slope and peat thickness on the water regime of a
forested gley soil.
Soil Use and Management **8** (2), 105–111.

Mechanics of wind damage

- Achim, A., Nicoll, B., Mochan, S. and Gardiner, B. (2003).
Wind stability of trees on slopes. In: *Wind Effects on Trees. Proceedings of the international conference*, eds B. Ruck, C. Kottmeier, C. Mattheck, C. Quine and G. Wilhel.
University of Karlsruhe, Germany, September 16–18, 2003, 231–238.
- Blackburn, P. and Petty, J. A. (1988).
Theoretical calculations of the influence of spacing on stand stability.
Forestry **61**, 235–244.
- Blackburn, P., Petty, J. A. and Miller, K. F. (1988).
An assessment of the static and dynamic factors involved in windthrow.
Forestry **61**, 29–43.
- Gardiner, B. A. (1989).
Mechanical characteristics of Sitka spruce.
Forestry Commission Occasional Paper 24.
Forestry Commission, Edinburgh.
- Gardiner, B. A. (1992).
Mathematical modelling of the static and dynamic characteristics of plantation trees. In: *Mathematical modelling of Forest Ecosystems*, eds J. Franke and A. Roeder, Sauerländers Verlag.
Frankfurt am Main, 40–61.
- Gardiner, B. A., Stacey, G. R., Belcher, R. E. and Wood, C. J. (1997).
Field and wind-tunnel assessments of the implications of respacing and thinning on tree stability.
Forestry **70**, 233–252.
- Holbo, H. R., Corbett, T. C. and Horton, P. J. (1980).
Aeromechanical behaviour of selected Douglas-fir.
Agricultural Meteorology **21**, 81–91.
- Mattheck, C. (1991).
Trees: The mechanical design.
Springer-Verlag, pp 121.
- Mayer, H. (1987).
Wind-induced tree sways.
Trees **1**, 195–206.
- Mayhead, G. J. (1973).
Some drag coefficients for British forest tree derived from wind tunnel studies.
Agricultural Meteorology **12**, 123–130.
- Mayhead, G. J., Gardiner, J. B. H. and Durrant, D. W. (1975).
Physical properties of conifers in relation to plantation stability.
Unpublished report, Forestry Commission, Edinburgh.

References & Further Reading

Milne, R. and Blackburn, P. (1989).
The elasticity and vertical distribution of stress within stems of *Picea sitchensis*.
Tree Physiology **5**, 195–205.

Milne, R. (1991).
Dynamics of swaying of *Picea sitchensis*.
Tree Physiology **9**, 383–399.

Monteith, J. L. (1975).
Vegetation and the atmosphere, Vol 1: Principles.
Academic Press, pp 278.

Monteith, J. L. (1976).
Vegetation and the atmosphere, Vol 1: Case Studies.
Academic Press, pp 439.

Nicoll, B.C., Achim, A., Mochan, S. and Gardiner, B.A. (2005).
Does steep terrain influence tree stability? A field investigation.
Canadian Journal of Forest Research **35**, 2360–2367.

Petty, J. A. and Worrell, R. (1981).
Stability of coniferous tree stems in relation to damage by snow.
Forestry **54**, 115–128.

Petty, J. A. and Swain, C. (1985).
Factors influencing stem breakage of conifers in high winds.
Forestry **58**, 75–84.

Ray, D. and Nicoll, B. C. (1998).
The effect of soil water-table depth on root-plate development and stability
of Sitka spruce.
Forestry **71**, 169–182.

Root anchorage

Anderson, C. J., Coutts, M. P., Ritchie, R. M. and Campbell, D. J. (1989).
Root extraction force measurements for Sitka spruce.
Forestry **62**, 127–137.

Blackwell, P. G., Rennolls, K. and Coutts, M. P. (1990).
A root anchorage model for shallowly rooted Sitka spruce.
Forestry **63**, 73–91.

Booth, T. C. and Mayhead, G. J. (1972).
Tree pulling. *Forestry Commission Report on Forest Research*, 65–66.

Coutts, M. P. (1983).
Root architecture and tree stability.
Plant and Soil **71**, 171–188.

Coutts, M. P. (1986).

Components of tree stability in Sitka spruce on peaty gley soil.

Forestry **59**, 173–197.

Coutts, M. P. Nielsen, C. N. and Nicoll, B. C. (2000).

The development of symmetry, rigidity and anchorage in the structural root system of conifers. In: *The supporting roots of trees and woody plants: Form function and physiology*, ed A. Stokes.

Kluwer Academic Publishers, 3–17.

Fraser, A. I. and Gardiner, J. B. H. (1967).

Rooting and stability in Sitka spruce.

Forestry Commission Bulletin 40.

HMSO, London.

Gibbs, J. N. and Greig, D. A. (1990).

Survey of parkland trees after the great storm of 16 October, 1987.

Arboricultural Journal **14**, 321–347.

Nicoll, B. C., Easton, E. P., Milner, A. D., Walker, C. and Coutts, M. P. (1995).

Wind stability factors in tree selection: distribution of biomass within root systems of Sitka spruce clones.

In: *Wind and trees*, eds M. P. Coutts and J. Grace, J.

Cambridge University Press, Cambridge, 276–292.

Nicoll, B.C., Gardiner, B.A., Rayner, B. and Peace, A.J. (2006).

Anchorage of coniferous trees in relation to species soil type and rooting depth.

Canadian Journal of Forest Research **36**, 1871–1883.

Quine, C. P. and Burnand, A. C. (1991).

Early growth and root architecture of Sitka spruce in relation to cultivation of a peaty ironpan afforestation site.

Scottish Forestry **45** (3), 175–182.

Quine, C. P., Burnand, A., Coutts, M. P. and Reynard, B. (1991).

Effects of mounds and stumps on the root architecture of Sitka spruce on a peaty gley restocking site.

Forestry **64** (4), 385–401.

Adaptive growth

Nicoll, B. C. and Ray, D. (1996).

Adaptive growth of tree root systems in response to wind action and site conditions.

Tree Physiology **16**, 891–898.

Ruel, J. C., Larouche, C. and Achim, A. (2003).

Changes in root morphology after precommercial thinning in balsam fir stands.

Canadian Journal of Forest Research **33**, 2452–2459.

References & Further Reading

Telewski, F. W. and Beals, W. J. (1995).
Wind induced physiological and developmental responses in trees.
In: *Wind and trees*, eds M. P. Coutts & J. Grace.
Cambridge University Press, Cambridge, pp 237–263.

Stokes, A., Nicoll, B. C., Coutts, M. P. and Fitter, A. H. (1997).
Responses of young Sitka spruce clones to mechanical perturbation and
nutrition: effects on biomass allocation, root development and resistance to
bending.
Canadian Journal of Forest Research **27**, 7–15.

Risk

Coutts, M. P. (1983).
Development of the structural root system of Sitka spruce.
Forestry **56**, 1–16.

Coutts, M. P., Walker, C. W. and Burnand, A. C. (1990).
Effect of establishment method on root form of lodgepole pine and Sitka
spruce and on the production of adventitious roots.
Forestry **63**, 143–159.

Gardiner, B. A., Peltola, H. and Kellomäki, S. (2000).
Comparison of two models for predicting the critical wind speeds required
to damage coniferous trees.
Ecological Modelling **129** (1), 1–23.

Gardiner, B. A. and Quine, C. P. (2003).
Wind damage to forests.
Biomimetics **2**, 139–147.

Gardiner, B. A., Suarez, J. and Quine, C. P. (2003).
Development of GIS based wind risk system for British forestry. In: *Wind
effects on trees. Proceedings of the international conference*, B. Ruck, C.
Kottmeier, C. Mattheck, C. Quine and G. Wilhelm (2003).
University of Karlsruhe, Germany, September 16–18, 2003, 145–150.

Gardiner, B. A., Peltola, H. and Kellomäki, S. (2003).
Comparison of two models for predicting the critical wind speeds required
to damage coniferous trees.
Ecological Modelling **129** (1), 1–23.

Miller, K. F. (1985).
Windthrow Hazard Classification.
Forestry Commission Leaflet 85.
HMSO, London.

Quine, C. P. (1991).
Recent storm damage to trees and woodlands in southern Britain.
In: *Research for practical arboriculture*, ed S. J. Hodge. Forestry Commission
Bulletin 97.
HMSO, London, 83–89.

Quine, C. P. (1995).
Assessing the risk of wind damage; practice and pitfalls.
In: *Wind and trees*, eds M. P. Coutts and J. Grace.
Cambridge University Press, Cambridge, 379–403.

Quine, C. P. and Wright, J. A. (1993).
The effects of revised windiness scores on the calculation and distribution of windthrow hazard class.
Forestry Commission Research Information Note 231.
Forestry Commission, Edinburgh.

Smith, K. (1992).
Environmental hazards; assessing risk and reducing disaster.
Routledge, London.

Silviculture

Hibberd, B. J. (1991).
Forestry Practice.
Forestry Commission Handbook 6.
HMSO, London.

Mason, W. L. (2002).
Are irregular stands more windfirm?
Forestry **75** (4), 347–355.

Quine, C. P. and Miller, K. F. (1991).
Windthrow – a factor influencing the choice of silvicultural systems. In:
Silvicultural systems, ed P. Gordon.
ICF, Edinburgh, 71–81.

Quine, C. P. and Gardiner, B. A. (1992).
Incorporating the threat of windthrow into forest design plans.
Forestry Commission Research Information Note 220.
Forest Research, Farnham.

Savill, P. S. (1983).
Silviculture in windy climates.
Forestry Abstracts **44**, 473–488.

Somerville, A. (1980).
Wind stability: forest layout and silviculture.
New Zealand Journal of Forestry Science **10**, 476–501.

Tree measurement and yield forecasting

Edwards, P. N. and Christie, J. M. (1981).
Yield models for forest management.
Forestry Commission Booklet 48.
Forestry Commission, Edinburgh.

References & Further Reading

Hamilton, G. J. (1975).
Forest mensuration handbook.
Forestry Commission Booklet 39.
HMSO, London.